

A. Shelf 10


LIBRARY

OF THE

U. S. Department of Agriculture.

Class 411
8-159 Sh2

v. 16



JACOBO PARKINSONO ARMIGERO,
IN
MUSEI CELEBERRIMI LEVERIANI
ADMINISTRATIONE
FACILI ET INGENUA LIBERALITATE
COMMENDATISSIMO,
DECIMUM SEXTUM

HUNC
NATURÆ VIVARIII
FASCICULUM

D. D. D.


GEORGIUS SHAW.

E. NODDER.

TO
JAMES PARKINSON ESQUIRE,
THE WORTHY AND LIBERAL PROPRIETOR
OF THE CELEBRATED
LEVERIAN MUSEUM,
THIS SIXTEENTH VOLUME OF
THE
NATURALIST'S MISCELLANY
IS WITH MUCH REGARD INSCRIBED
BY
GEORGE SHAW.
E. NODDER.

LIBRARY
U. S. DEPARTMENT
OF AGRICULTURE

LANIUS GUTTURALIS.



CHARACTER GENERICUS.

Rostrum rectiusculum, dente utrinque versus apicem, basi nudum.

Lingua lacera.

Lin. Syst. Nat. p. 134.

CHARACTER SPECIFICUS, &c.

LANIUS olivaceus, gula coccinea, lunula pectorali nigra, fronte humerisque flavis.

LANIUS GUTTURALIS. L. viridi-olivaceus, mento, gula, infero pectore et ano coccineis; fronte humerisque flavis, cum linea oculari in lata lunula pectorali transversa protensa.

DAUDIN. *Ann. Mus. d'Hist. Nat.*

In regionibus Africanis calidioribus generatur *Lanius gutturalis*, eadem magnitudine qua *Lanius Excubitor* seu vulgaris. Sedem præcipue seligere dicitur in excelsis arborum cacuminibus, et in hoc plerisque sui generis esse dissimilis quod baccis vescatur.

THE
RED-THROATED BUTCHER-BIRD.

GENERIC CHARACTER.

Bill straitish, with a tooth or small process on each side near the tip.

Tongue jagged.

SPECIFIC CHARACTER, &c.

Olive BUTCHER-BIRD, with scarlet throat, black pectoral crescent, and yellow front and shoulders.

Pie-Grièche à gorge rouge.


DAUDIN. *Ann. Mus. d'Hist. Nat.*

This bird is a native of the warmer parts of Africa, and is equal in size to the great grey or common Butcher-Bird. It is said chiefly to frequent the tops of lofty trees, and, contrary to most of the genus, to feed on various kinds of berries.



R. W. Dole del.

APLYSIA PUNCTATA.



CHARACTER GENERICUS.

Corpus repens obvelatum membranis reflexis,
clypeo dorsali pulmones obtegente.

Foramen laterale dextrum.

Anus supra extremitatem dorsi.

Tentacula quatuor, anterieus sita.

CHARACTER SPECIFICUS, &c.

APLYSIA purpurea albido-punctata.

LAPLYSIA PUNCTATA.

CUVIER. *Ann. d'Hist. Nat.*

Aplysiam quam depinximus primus descripsisse videtur celeberrimus Cuvierus utpote speciem genuinam, et a reliquis vere sejunctam. Narrat ille non raro latere eam sub saxis lapidibusque littoris Massiliensis, vel ipso linace vulgari seu hortensi tardius incedere, vescique parvulis testaceis, et reliquis

quis ejusmodi. Si quis eam turbaverit seu contrec-
taverit, liquorem purpureum e corporis meatibus
emittere solet, aquasque propius circumfusas con-
colores reddere. Creditur Aplysiam punctatam pro-
lis esse fœcundam, pullosque spatio bimestri plene
adolevisse.



THE
SPECKLED APLYSIA.


GENERIC CHARACTER.

SPECIFIC CHARACTER.

The present species of *Aplysia* appears to have been first distinguished as a separate species by the celebrated Mons. Cuvier, who assures us that it is very

very common about the coasts of Marseilles, generally lying concealed beneath rocks or stones, moving in a slower manner than even the common garden slug, and feeding on very small testacea, &c. When disturbed or handled it discharges from its pores a purple juice, discolouring the water to some small distance round. It is supposed to be a prolific animal, and the young are said to arrive at their full size in the space of two months at farthest.

OSTREA JACOBÆA.



CHARACTER GENERICUS.

Animal Tethys.

Testa bivalvis, inæquivalvis, subaurita.

Cardo edentulus fossula cava ovata striisque lateralibus transversis.

CHARACTER SPECIFICUS, &c.

OSTREA JACOBÆA. O. testa inæquivalvi radiis quatuordecim angulatis longitudinaliter striatis.

Gualt. test. t. 99. f. B.

Lesser testacoth. f. 139.


In oris Europæis sæpius reperitur *Ostrea Jacobæa*, coloribus non raro varians, *Ostrea maxima*, ut plurimum, paulo minor.

Hinge toothless, with an ovate fossule and transverse lateral streaks.

SCALLOP with fourteen angular, longitudinally-striped rays.

Penn. Brit. Zool. pl. 60. f. 62.

The Jacobæan Scallop is sufficiently common about the European coasts, often varying in colour, and generally rather smaller than the common Scallop.



R.P.N. Del^t Sculp^t

PHALÆNA APOLLONIA.



CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ. (Volatu nocturno.)

Lin. Syst. Nat. p. 808.

CHARACTER SPECIFICUS, &c.

PHALÆNA alba, alis omnibus macula ocellata fasciisque transversis nigris luteo marginatis.

PHALÆNA APOLLONIA.

Cram. t. 250. f. A.

Attaci.

Africam Australem incolit pulchra hæc phalæna, magnitudine vera in tabula depicta.



R.P.N. Del. Sculp.

APOLLONIA.



GENERIC CHARACTER.

Antennæ setaceous, decreasing in size from base to point.

Wings (when at rest) generally deflected.

Flight nocturnal.

SPECIFIC CHARACTER.

White *Phalæna* with all the wings marked by a black ocellated spot and transverse black bands edged with yellow.

This beautiful *Phalæna* is a native of the southern parts of Africa, and is represented on the plate in its natural size.

PAVO SPICIFER.



CHARACTER GENERICUS.

Caput pennis revolutis tectum.

Pennæ uropygii elongatæ, ocellatæ.

Lin. Syst. Nat. p. 267.

CHARACTER SPECIFICUS, &c.

PAVO capite cæruleo, pennis cristæ lanceolatis erectis, fascia utrinque suboculari lutea.

PAVO muticus. P. capite crista subulata, calcaribus nullis?

PAVO japonensis.

Lin. Syst. Nat. p. 268.

Aldrov. orn. 2. p. 35.

Eandem sibi patriam vindicat pulcherrima hæc avis quam et pavo vulgaris; Indiam scilicet, ibique præcipue regiones montanas. Moribus modoque vivendi pavoni vulgari similis esse dicitur, a quo etiam verisimile est sæpius non fuisse discretam. Speciem tamen revera esse diversam satis probat

B

peculiaris

peculiaris cristæ conformatio, nec non macula saturatim flava utrinque sub oculis sita. Si idem sit *pavo muticus* Linnæi, (*spicifere* Buffoni) atque hæc nostra avis; vel in illo describendo evidenter erratum est, vel variare solent colores et alia nonnulla; Linnæus enim, et Aldroyandus, (qui primus descriptor avem quam depictam adeptus est, repræsentavit,) calcaribus carentem ostendunt; quæ in icone unde deducta est nostra similitudo, admodum conspicua sunt: hanc autem effigiem ipsius avis viventis expressit pictor Indicus, eamque nobiscum perhumaniter communicavit Dominus Fleming, chirurgus Anglicus qui prope Calcuttam in Bengala multos annos versatus est.



THE
SPIKE-CRESTED PEACOCK.



GENERIC CHARACTER.

Head covered with revolute feathers.

Uropygial feathers elongated and ocellated.

SPECIFIC CHARACTER.

PEACOCK with blue head, upright lanceolate crest, and yellow band beneath each eye.

The beautiful bird represented on the present plate is, like the common Peacock, a native of India, chiefly inhabiting mountainous regions. It is distinguished as a species by the peculiar structure of its crest, as well as by the bright-yellow patch beneath each eye. In its manners it is said to resemble the common peacock, with which it has probably been often confounded. If it be the bird intended by the *Pavo muticus* of Linnæus, (the *Spicifere* of Buffon,) it is evident either that the description of that species is erroneous, or that the bird is subject

to vary in colour and other circumstances; since Linnæus and Aldrovandus, (who first described it from a painting,) represent it as destitute of spurs, which are conspicuous in the drawing from which the present figure was copied, and which was taken from the life in India, and politely communicated by T. Fleming, Esq. several years resident in the neighbourhood of Calcutta in Bengal.

CANCER MANTIS.



CHARACTER GENERICUS.

Pedes octo (rarius sex aut decem;) insuper manus duæ chelatae.

Oculi duo, distantes, plurimis pedunculati; elongati, mobiles.

Cauda articulata, inermis.

CHARACTER SPECIFICUS, &c.

CANCER MANTIS. *C. macrourus* articularis, manibus adactylis, compressis, falcatis, serrato-dentatis.

Lin. Syst. Nat. p. 1054.

Squilla MANTIS.

Fabr. sp. ins. 1. p. 514.

CANCER arenarius.

Rumph. mus. t. 3. *f.* 2.

Nomen adeptus est *Cancer Mantis* ab exilitate et gracilitate corporis; qua in re accedit quodammodo
ad

ad similitudinem insectorum quæ complectitur genus Mantis. In mari innascitur mediterraneo, repertus præcipue apud oras Italicas.

In systemate Fabriciano non satis ad normam redacto datur huic speciei genus distinctum nomine Squillæ.

De novis plurimis generibus nuperrime institutis liceat mihi dicere, me iudice, obscurari ab iis insectorum historiam magis quam elucidari, quique antea satis clarus erat et facilis investigationis modus, eum incertum sæpius et difficilem reddi.

Major sæpe est Cancer Mantis quam qui in tabula ostenditur.



THE
LONG-BODIED CRAB.



GENERIC CHARACTER.

Legs generally eight, (in some species six or ten,) besides two claspers or chelated arms.

Eyes two, commonly distant; footstalked, moveable.

Tail jointed.

SPECIFIC CHARACTER, &c.

Long-tailed many-jointed CRAB, with fingerless, compressed, falcated, tooth-serrated chelæ.

The Cancer Mantis, so named from its long and slender habit, in which it in some measure resembles the insects of the genus *Mantis*, is a native of the mediterranean sea, and is principally observed about the Italian coasts.

In the singular and irregular system of Fabricius, this species ranks under a distinct genus entitled *Squilla*.

I may

I may here be permitted to observe, that of many new genera which the spirit of modern entomologists has endeavoured to establish it may with truth be affirmed that they have rather tended to obscure than elucidate the history of insects, by rendering that investigation troublesome and uncertain which was before sufficiently easy and clear.

The Cancer Mantis is often found of a much larger size than represented on the present plate.

MEDUSA NUMMULARIA.



CHARACTER GENERICUS.

Corpus gelatinosum, orbiculatum, depressam.

Os subtus, centrale.

Lin. Syst. Nat. p. 1096.

CHARACTER SPECIFICUS, &c.

MEDUSA orbicularis albida subpellucida radiata, subtus appendicibus tribus cæruleis.

Le PORPITE appendiculée.

Nouv. Dict. d'Hist. Nat.


Pulchram hanc Medusam, marium septentrionalium incolam, juxta magnitudinem naturalem depinximus. Radit sæpius hæc species summas aquas catervatim congregata. Habitu seu facie generali simillima est Medusæ Porpitæ Linnæi, quam inter et hanc nostram sæpius nullum discrimen fecisse videntur descriptores.

NUMMULAR MEDUSA.

GENERIC CHARACTER


643

Body flattened, or depressed.
Abdomen broadly, or



SPECIFIC CHARACTER

Whitish, orbicular, subpellucid, radiated. 15-
DUSA with
width



The beautiful Medusa represented in its natural
size on the present plate is a native of the northern
ocean, where it is often observed. The name
R. P. N. Del. et Sculp.
has swimming on the surface of the water. In its
general habit it bears the greatest resemblance to
the Medusa *Physalia* of Linnaeus, with which it ap-
pears to have been generally confounded.

THE
NUMMULAR MEDUSA.



GENERIC CHARACTER.

Body gelatinous, orbicular, depressed.
Mouth beneath, central.

SPECIFIC CHARACTER.

Whitish, orbicular, subpellucid, radiated MEDUSA, with three blue appendices beneath.

The beautiful Medusa represented in its natural size on the present plate is a native of the northern seas, where it is often observed in considerable numbers, swimming on the surface of the water. In its general habit it bears the greatest resemblance to the Medusa Porpita of Linnæus, with which it appears to have been generally confounded.

PHALÆNA FABIA.



CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ. (volatu nocturno.)

Lin. Syst. Nat. p. 808.

CHARACTER SPECIFICUS, &c.

PHALÆNA alis rotundatis luteis, macula marginali lineisque transversis undulatis fuscis.

PHALÆNA Fabia.

Cram. t. 256. *f.* B.

Varias Indiæ partes incolit Phalæna Fabia, cujus veram magnitudinem ostendit tabula.



W. V. Decker Del. J. G. L. Sculp.

FABIA.



GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to point.

Wings (when at rest) generally deflected.

Flight nocturnal.

SPECIFIC CHARACTER.

MOTH with deep-yellow rounded wings, with brown marginal spot and transverse undulated lines.

PHALÆNA FABIA.

Cram. t. 250. f. B.

The *Phalæna Fabia* is a native of many parts of India, and is represented on the plate in its natural size.

LANIUS JOCOSUS.



CHARACTER GENERICUS.

Rostrum rectiusculum, dente utrinque versus
apicem, basi nudum.

Lingua lacera.

Lin. Syst. Nat. p. 134.

CHARACTER SPECIFICUS, &c.

LANIUS fuscus capite nigro, macula subocu-
lari crissoque coccineis.


LANIUS JOCOSUS. L. cauda rotundata, cor-
pore griseo, palpebra inferiore purpurea,
ano sanguineo.

Lin. Syst. Nat. p. 138.

MERULA sinensis cristata minor.

Briss. 2. p. 255. t. 21. f. 2.

Avem Sinæ incolam magnitudine deminuta repræ-
sentavimus, inter pulcherrimas sui generis species
numerandam, ingenio præditam agili alacrique, et
apto ad imitationem,





THE
RED-VENTED SHRIKE.



GENERIC CHARACTER.

Bill straitish, with a tooth or small process on each side near the tip.

Tongue jagged.

SPECIFIC CHARACTER, &c.

Brown SHRIKE, with black head, red vent, and red spot beneath the eyes.

JOCOSE SHRIKE.

Lath. Syn. 1. p. 175.

La petit Merle hupé de la Chine.

Briss. orn. 2. t. 21. f. 2.

This bird, one of the most beautiful of the genus, is a native of China, and is represented smaller than the natural size : it is remarkable for the liveliness and docility of its disposition.

PHALÆNA TARQUINIA.



CHARACTER GENERIC.

Antennæ setacæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ (volatu nocturno.)

Lin. Syst. Nat. p. 808.


CHARACTER SPECIFICUS, &c.

PHALÆNA alis superioribus nigris, linea media furcata alba, inferioribus fulvis.

PHALÆNA TARQUINIA.

Cram. t. 4. f. A.

Americam australem incolit Phalæna Tarquinia
cujus veram magnitudinem ostendit tabula.



646



R.P. Nodder

TARQUINIA.



GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to tip.

Wings (when sitting) generally deflected.
(Flight nocturnal.)

SPECIFIC CHARACTER, &c.

MOTH with the upper wings black, marked by a forked middle line; the lower wings fulvous.

TARQUINIA.

Cram. pap. 1. *pl.* 4. f. A.

This elegant species is a native of South-America, and is represented on the plate in its natural size.

TARQUINIA

TARQUINIA

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

Characteristics of the species

LUCERNARIA QUADRILOBA.



CHARACTER GENERICUS.

Corpus gelatinosum, rugosum, ramosum.

Os inferum.

Lin. Syst. Nat. Gmel. p. 3151.

CHARACTER SPECIFICUS, &c.

LUCERNARIA rubra, brachiis quatuor apice tentaculatis.



LUCERNARIA quadricornis. L. Corpore elongato tortili, brachiis quatuor dichotomis, apice tentaculatis.

Lin. Syst. Nat. Gmel. p. 3151.

Müll. Zool. Dan. t. 39. f. 1.—6.

Mirum hoc Zoophyton, cui cum hydrys arctissima videtur esse affinitas, ad genus ab iis separatum, et nomine *Lucernariæ* distinctum, primus retulit celeberrimus Müllerus in opere cui titulus *Zoologia Danica*. Maria incolit septentrionalia, repertum præcipue

cipue circa oras Danicas et Norvegianas; adhæretque
fucis, aliisque multis quæ in oceano nascuntur. Nec
ab ipsis Britannis omnino ignoratur, cum paucis ab-
hinc annis perpulchrum detexerit specimen prope
littus Hibernicum Dominus Templetonus Linnæanæ
societatis Londinensis socius. Exhibet tabula na-
turalem magnitudinem.



R. N. D. 100, 100

3

THE
FOUR-LOBED LUCERNARIA.



GENERIC CHARACTER.

Body gelatinous, wrinkled, branched.

Mouth beneath.

SPECIFIC CHARACTER, &c.

Red LUCERNARIA, with four arms tentaculated at the tip.

Müll. Zool. Dan. t. 39. f. 1.—6.


This highly curious Zoophyte, allied in the strongest manner to the Polypes, seems to have been first considered as forming a distinct genus by the celebrated Müller in his *Zoologia Danica*, under the title of Lucernaria. It is a native of the northern seas, and has been chiefly observed about the shores of Denmark and Norway, adhering to fuci and other marine substances; nor is it altogether a stranger to the British isles, a very fine specimen having been discovered

discovered some years past on the coast of Ireland
by the ingenious Mr. Templeton, Fellow of the Lin-
naean Society. The plate represents it in its natural
size.

CYPRÆA FRAGILIS

ET

CYPRÆA ARABICA.



CHARACTER GENERICUS.

Animal Limax.

Testa univalvis, involuta, subovata, obtusa,
lævis.

Apertura utrinque effusa, linearis, utrinque den-
tata, longitudinalis.

Lin. Syst. Nat. p. 1172.

CHARACTER SPECIFICUS, &c.

CYPRÆA FRAGILIS. C. testa turbinata ovata
glauca, testaceo-undata subfasciata.

Lin. Syst. Nat. p. 1175.

Mus. Lud. Ulric. 570. n. 188.

Gualt. test. t. 16. f. q.

Habitat in mari mediterraneo.

CHARACTER SPECIFICUS, &c.

CYPRÆA ARABICA. C. testa subturbinata
characteribus inscripta, macula longitudi-
nali simplici.

Lin. Syst. Nat. p. 1173.

Percellana literata s. ARABICA.

Rumph. mus. t. 38. f. m.

List. Conch. t. 658. f. 3?

Gualt. test. t. 16. f. v.

Habitat in mari Indico.

648



R. P. Nodder D. L. et. Sculp.

Aperture linear, longitudinal, toothed on both sides.

BRITTLE COWRY. Glaucous-brown Cowry
with yellowish-brown bands and streaks.

Knorr. vergn. 5. t. 18. f. 2.

Native of the Mediterranean sea.

SPECIFIC CHARACTER, &c.

ARABIAN COWRY. Subturbinated Cowry
with characteriform variegations and violet-
coloured mouth.

List. Conch. t. 658. f. 3.

Native of the Indian seas.

MOTACILLA ROSSICA.



CHARACTER GENERICUS.

Rostrum subulatum, rectum; mandibulis sub-
æqualibus.

Nares obovatæ.

Lingua lacero-emarginata.

Lin. Syst. Nat. p. 328.

CHARACTER SPECIFICUS, &c.

MOTACILLA grisea, capite nigro, collo utrin-
que albo, pectore abdomineque croceis.

MOTACILLA rubicola. var. β .

Lin. Syst. Nat. Gmel.

MOTACILLA superne nigricans, torque albo
interrupto, pectore atque abdomine supe-
riore croceis.

Lepechin Act. Petrop. 15. t. 25. f. 3.

In Russiæ et Sibiriae betuletis atque locis paludo-
sis præcipue conspicitur Motacilla Rossica, a Domi-
no Lepechin in Actis Petropolitanis ita descripta.

“ Rostrum tenue nigrum, mandibula superior
paulo longior apice incurva uti in congeneribus:

vertex capitis, genæ, gula atque collum antèrius atra, nucha quoque insignitur nigredine a capillitio ad dorsum producta; partes laterales colli albæ; qui color etiam summa hypochondriorum tenet: pectus atque abdomen crocea, sed in abdomine croceus color magis magisque diluitur ita ut ad pedes albidus sit: dorsum nigricans margine pennarum parum rufescente: scapulæ alarum nivæ, rectrices anteriores nigræ apicibus ex albido rufescentibus: remiges majores fuscæ, minores nigricantes, omnes margine vexilli interni ad dimidium albo: uropygium utrinque niveum: rectrices duodecim æquales nigræ exceptis utrinque externis, quarum margo vexilli externi albicat: pedes unguesque nigri. Fæmina supra fusca marginibus pennarum rufescentibus, macula alarum candida, gutture sordide albo, pectore atque abdomine dilute rufescente: remiges rectricesque prouti in mare."



R. P. N. Del. et Sculp.

Tongue jagged, or lacerated towards the tip.

Lepech. Act. Petr. 15. p. 439.

D 2

black, which colour also reaches from the nape to the beginning of the back: the sides of the neck, and upper parts of the sides are white: the breast and abdomen orange, the colour gradually growing paler as it approaches the legs, where it becomes whitish: the back is blackish, the feathers being slightly rufous on their edges: the scapulars are white, the smaller coverts black, with rufous-white tips: the great wing-feathers brown, the smaller blackish, the inner edges of both being white for half their length: the rump and vent are white: the twelve tail-feathers are of equal length, and black, except the outside ones, which are whitish on the inner edges: the legs and claws are black. The female is brown above, with the feathers slightly edged with rufous: on each wing is a white patch: the throat is of a dull white: the breast and abdomen slightly rufous: wings and the tail similar to those of the male bird.

PAPILIO AMPHRYSIUS.



CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes ;
(volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis denticulatis nigris, versus apices flavo-substriatis, inferioribus flavis nigro-marginatis.

PAPILIO alis concoloribus nigris, anterioribus maculis, posterioribus disco flavis.

Lin. Syst. Nat. Gmel. p. 2230.

Eq. Tr.

Merian Ins. Surin. t. 72 ?

PAPILIO AMPHRYSIUS.

Cram. t. 219. f. A.

Affinis P. Helenæ.

Insulas Javam atque Amboinam præcipue incolit formosa hæc papilionis species, magnitudine vera in tabula depicta.



R.P. Nodder. Del: et. Sculp:!

AMPHRYSIUS.



GENERIC CHARACTER.

Antennæ generally thickening towards the extremities, and terminating in a clavated tip.

Wings (when sitting) erect, and meeting upwards.

(Flight diurnal.)

SPECIFIC CHARACTER, &c.

BUTTERFLY with denticulated wings; the upper pair black, with obscure yellow streaks towards the tips; the lower yellow, bordered with black.

Jablonsk. pap. 1. t. 1. f. 3.

This beautiful Butterfly is principally found in the islands of Java and Amboina, and is represented in its natural size on the annexed plate.

SPONDYLUS GEDDERTUS

CHARACTER GEDDERTUS

CHARACTER GEDDERTUS

Animal talipes
Tacta irregularis rigida
Corde dentibus duobus recurvis cum forami-
nulo intermedio
Linn. Syst. Nat. p. 1136

CHARACTER GEDDERTUS

SPONDYLUS testa subnulla spinosa
Linn. Syst. Nat. p. 1136
Gmel. test. non. Nat. p. 1136
Bonn. test. non. Nat. p. 1136
Reg. test. non. Nat. p. 1136

Ad hoc inter se generis spatium similitudine com-
municant. Spondyli et Cyprii generis, quia ambo ad
genus latere possunt, etiam si ut prius non du-
bitat, non possit de nonnullis speciebus, sicut et hoc
ad ad illud genus referre. Hoc tamen insignis
est ad distinctionem, quia Cyprii spondylus se-
cundum; nullo namque valvula interiore, qui quod
abscessus ad abscissus superficem habet complan-
tate. Species praecipuae, quam deplurimum spergit
se in innumeras varietates quoad colorem et magni-

SPONDYLUS GÆDEROPUS.



CHARACTER GENERICUS.

Animal Tethys.

Testa inæquivalvis, rigida.

Cardo dentibus duobus recurvis, cum foraminulo intermedio.

Lin. Syst. Nat. p. 1136.

CHARACTER SPECIFICUS, &c.

SPONDYLUS testa subaurita spinosa.

Lin. Syst. Nat. p. 1136.

Gualt. test. 100. 101. f. A. B.

Rumph. mus. t. 47. f. E.

Regenf. conch. 16. t. 4. f. 48.

Adeo inter se generali quadam similitudine commiscentur *Spondyli* et *Chamæ* genera, (quæ ambo ad genus *Ostreæ* proxime accedunt) ut primo visu dubitari fere possit de nonnullis speciebus, sintne ad hoc an ad illud genus referendæ. Nota tamen insignis est ad distinctionem, quæ a *Chamis* *Spondylos* secernit; umbo nempe valvulæ inferioris, qui quasi abscissus seu abrasus superficiem habet complanatam. Species præcipua, quam depinximus, spargit se in innumeras varietates quoad colorem et magni-

tudinem. E pulcherrimis speciminibus, quæ in Museo Leveriano asservantur, testam selegimus, cujus ad similitudinem figuram hanc nostram exprimi curavimus.



R.P.N. Del et Sculp

SUBAURICULATED SPONDYLUS.



GENERIC CHARACTER.

Animal allied to a Tethys.

Shell of unequal valves, strong.

Hinge with two recurved teeth and intermediate hollow.

SPECIFIC CHARACTER, &c.

SPONDYLUS with subauriculated, spinose shell.

Bonann. recr. t. 20. 21.

Seb. mus. 3. t. 88. f. 1—11.

The two genera of *Spondylus* and *Chama*, (both nearly allied to that of *Ostrea*,) bear so great a general resemblance to each other, that, in some species, it appears doubtful, at first sight, to which of the two the shell should be most properly referred. One great particularity however always distinguishes the shells of this genus from those of the genus *Chama*; viz. the aspect of the top of the inferior valve, which appears exactly as if sawn or cut away by an instrument, so as to leave a flat surface on that part. The



R. P. N. Del. et Sculp.

THE
BLACK-FINNED SPARUS.



GENERIC CHARACTER.

Teeth strong; *Front-Teeth* disposed in a single, double, treble, or quadruple row.

Grinders (in most species) convex, smooth, and disposed in ranges, forming a kind of pavement in the mouth.

Lips thick: *Gill-Covers* unarmed, smooth, scaly.

SPECIFIC CHARACTER, &c.

Purplish SPARUS, with yellowish-green abdomen, orbits of the eyes radiated by green streaks, and black fins.

Black-Finned SPARUS.

Gen. Zool. 4. p. 451.

The black-finned Sparus is seen chiefly in the Indian seas, and usually measures about twelve inches in length.

PSITTACUS MELANOTUS.



CHARACTER GENERICUS.

Rostrum aduncum ; mandibula superiore mobili, cera instructa.

Nares in rostri basi.

Lingua carnosae, obtusa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

CHARACTER SPECIFICUS, &c.

PSITTACUS macrourus viridis, dorso nigro, tectricibus sanguineis, uropygio cæruleo.

Psittacus Melanotus a nemine antea descriptus proculdubio habendus est in iis quos pulcherrimos alit Australasia. Psittaco *Pennantii* et illo qui *eximius* dicitur paulo minor colores jactat egregie vividos et elegantes.

THE
BLACK-BACKED PARRAKEET.

GENERIC CHARACTER.

Bill hooked: upper mandible moveable, and furnished with a cere.

Nostrils in the base of the bill.

Tongue fleshy obtuse, generally entire.

Feet scansorial.

SPECIFIC CHARACTER, &c.

Long-tailed green PARRAKEET, with black back, scarlet wing-coverts, and blue rump.

The Parrakeet represented on the present plate is unquestionably one of the most beautiful that have yet been discovered in the regions of Australasia, and is a species hitherto undescribed. Its size is somewhat smaller than that of the Pennantian and nonpareil parrakeets, and its colours in the highest degree vivid and elegant.

SCOMBER PLUMIERI.



CHARACTER GENERICUS.

Corpus oblongum, læve, linea laterali interdum carinatum.

Pinnulæ sæpius supra infraque versus caudam.

CHARACTER SPECIFICUS, &c.

SCOMBER viridi-cæruleus, subtus argenteus, squamis magnis, pinnis pectoralibus lanceolatis, linea laterali rotundato-loricata.

SCOMBER PLUMIERI. S. squamis magnis, pinnis flavis.

Bloch. t. 344.

Maria incolit Americana Scomber Plumieri, longitudine, ut plurimum pedali.

657



RPN Del Sulp

THE
PLUMIERIAN MACKREL.



GENERIC CHARACTER.

Body oblong, smooth, sometimes carinated by the lateral line.

Finlets (in most species) above and below, towards the tail.

SPECIFIC CHARACTER, &c.

Blue-green MACKREL, silvery beneath, with large scales, lanceolate pectoral fins, and roundly-mailed lateral line.

The Plumierian Mackrel is an inhabitant of the American seas, and usually measures about twelve inches in length.

CANCER SCYLLARUS.



CHARACTER GENERICUS.

Pedes octo (rarius sex aut decem;) insuper
manus duæ chelatæ.

Oculi duo, distantes, plurimis pedunculati;
elongati, mobiles.

Cauda articulata, inermis.

CHARACTER SPECIFICUS, &c.

CANCER macrourus articularis, manibus adac-
tylis, ventricosis rectis angulatis introrsum
tridentatis.

Lin. Syst. Nat. p. 1054.

SQUILLA arenaria.

Seb. mus. 3. t. 20. f. 6.

Cancro *Manti*, quem in opere hoc nostro non
ita pridem repræsentavimus, licet longe minor sit
Cancer *Scyllarus*, in aliis tamen conjungi videtur
cum eo propinqua cognatione. In maribus enutri-
tur Indicis,



RPN

THE
OVAL-TAILED CRAB.



GENERIC CHARACTER.

Legs generally eight, (in some species six or ten,) besides two claspers or chelated arms.

Eyes two, commonly distant; footstalked, moveable.

Tail jointed.

SPECIFIC CHARACTER, &c.

Long-tailed many-jointed CRAB, with adactyle, ventricose, strait, angular arms, tridentated on the inner side.

Rumph. mus. t. 3. f. F.

This species, though of much smaller size, is very nearly allied in point of general resemblance to the Cancer Mantis lately figured in the present work, and is a native of the Indian seas.

PHALÆNA JANUS.



CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ. (Volatu nocturno.

Lin. Syst. Nat. p. 808.

CHARACTER SPECIFICUS, &c.

PHALÆNA alis primoribus fuscis griseo-fasciatis, posterioribus rubris macula media magna ocellata nigra.

PHALÆNA alis patulis, anticis variegatis, subtus ocello atro, posticis sanguineis ocello atro.

Fab. sp. ins. 2. p. 169.

Insectum depinximus amplum et venustum, cum *Phalæna Augusta*, de qua antea in hoc opere disseruimus, remota quadam cognatione conjunctum. In America Australi generatum sedem sibi præcipuam seligere dicitur in Surinamia. Ostenditur in tabula naturalis magnitudo.

657



JANUS.



GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to point.

Wings (when at rest) generally deflected.

Flight nocturnal.

SPECIFIC CHARACTER, &c.

PHALÆNA with the upper wings brown with grey bands, the lower red with a large black ocellated spot.

PHALÆNA JANUS.

Cram. pl. 64. f. A. B.

The large and elegant insect here exhibited, and which is in some degree allied to the *Phalæna Augusta*, before represented in the present work, is a native of South-America, and is said to be principally found in Surinam. The plate expresses it in its natural size.

PSITTACUS ZONARIUS.



CHARACTER GENERICUS.

Rostrum aduncum; mandibula superiore mobili, cera instructa.

Nares in rostri basi.

Lingua carnosae, obtusa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

CHARACTER SPECIFICUS.

PSITTACUS macrourus viridis, capite remigibusque nigris, torque nuchali fasciaque abdominali flavis.

De hac specie satis sit notare novam omnino esse nec antea descriptam. In Australasia generatur, magnitudine quasi Psittaci Tabuani.

ZONED PARROT.



THE
ZONED PARROT.



GENERIC CHARACTER.

Bill hooked: upper mandible moveable, and furnished with a cere.

Nostrils in the base of the bill.

Tongue fleshy, obtuse, generally entire.

Feet scansorial.

SPECIFIC CHARACTER.

Long-tailed green PARROT, with the head and long wing-feathers black, the collar on the nape and abdominal zone yellow.

Of this species it may be sufficient to observe that it is new and hitherto undescribed. Its size is that of the Tabuan Parrot.



PHALÆNA PAPHIA.



CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ. (Volatu nocturno.

Lin. Syst. Nat. p. 808.

CHARACTER SPECIFICUS, &c.

PHALÆNA pectinicornis elinguis flava, alis falcatis concoloribus ocello fenestratis.

Lin. Syst. Nat. p. 809.

PHALÆNA alis patentibus concoloribus flavis, strigis rufis ocelloque fenestrato.

Fabr. sp. ins. 2. p. 168.

Phalænam quam depinximus generant multæ Indicæ regiones; ibique larva ejus prægrandis et quasi bombycina nomine *Tusseh* distinguitur. Rhamni Jujubæ Linnæi (quam *Byer* nuncupant Hindostaniam incolentes) folia præcipue depascitur; et a Bengalensi populo copiose colligitur, ut
inde

indē deducatur sericum subfuscum, crassum, firmum et durabile, quo vestiri solent Brachmanes, et alii nonnulli religiosi. Involvitur chrysalis tela seu theca ovata, unde comparatur sericum, quam ab extremo ramulis affigit chorda valida e filamentis agglutinatiis contorta. Postquam novem menses transegerit chrysalis, nascitur phalæna, mense præcipue Julio; cujus color variat, interdum flavus, interdum fusco seu gilvo leviter suffusus. In tabula nostra exprimitur phalæna, nec non larva, magnitudine naturali.

Larvæ imaginem et notitiam debemus ingenioso Domino Roxburgh, qui de illa disseruit in septimo volumine actorum Societatis Linnæanæ.



Sp. 1st Ed. 1st. 1st.

PAPHIA.



GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to point.

Wings (when at rest) generally deflected.

Flight nocturnal.

SPECIFIC CHARACTER, &c.

Yellow PHALÆNA, with patulous subfalcated wings, marked by a rufous stripe and transparent ocellated central spot.

PHALÆNA Mylitta.

Drury ins. 2. t. 5. *Cram. t.* 146. 147.

The beautiful Phalæna here represented is a native of many parts of India, where its caterpillar, which grows to a large size, is known by the name of the Tusseh Silk-worm. It feeds principally on the leaves of the *Rhamnus Jujuba* of Linnæus, (called Byer among the Hindoos,) and is so plentiful in many

many parts of Bengal as to afford to the natives, who collect it for that purpose, a strong, dark-coloured, coarse silk, extremely durable, and serviceable for many articles of dress, being much worn by the Bramins and some other sects. The webs in which the animal changes into its chrysalis state, and which afford the silk above-mentioned, are of an oval shape, and fastened at one end to the twigs on which they are placed by a very strong cord of agglutinated filaments. The moth makes its appearance chiefly in the month of July, having lain nine months in its chrysalis state. In colour it varies, being sometimes more or less tinged with pale brown or buff, instead of clear pale yellow, and, as well as the caterpillar, is exhibited on the plate in its natural size. For the above history of the insect, with the figure of the larva or caterpillar, we are obliged to the ingenious Dr. William Roxburgh, whose description of the animal is inserted in the seventh volume of the Transactions of the Linnæan Society.

CORALLINA FLABELLUM.



CHARACTER GENERICUS.

Animal ? crescens habitu plantæ.

Stirps fixa. Rami articulati, ramulosi.

CHARACTER SPECIFICUS, &c.

CORALLINA FLABELLUM. C. stipite simplici incrustato, &c.

Lin. Syst. Nat. Gmel. p. 3842.

CORALLINA stipite simplici incrustato, ramis omnibus conglutinated, fronde flabelliformi incrustata subundulata.

Soland. et Ellis Zooph. p. 124. t. 24.

Rupibus maris Americani adhærescit elegans hæc Corallinæ species; interdum major quam in tabula depicta.



R. P. Nodder Del. et Sculp.

THE
FAN CORALLINE.

GENERIC CHARACTER.

Animal? growing in the form of a plant.

Stem fixed. Branches jointed and subdivided.

SPECIFIC CHARACTER.

Greenish-white CORALLINE, with simple stem, conglutinated branches, and fan-shaped subundulated frond.

Fan-shaped CORALLINE.

This elegant species of Coralline is found on the rocks of the American seas, and sometimes arrives at a larger size than represented on the plate.

MOTACILLA SUECICA.



CHARACTER GENERICUS.

Rostrum subulatum, rectum; mandibulis sub-
æqualibus.

Nares ovatæ.

Lingua lacero-emarginata.

Lin. Syst. Nat. p. 328.

CHARACTER SPECIFICUS, &c.

MOTACILLA fusca, pectore caudæque basi
ferrugineis, gula cærulea, abdomine albido.

MOTACILLA SUECICA. M. pectore ferru-
gineo fascia cærulea, rectricibus fuscis,
versus basin ferrugineis.

Lin. Syst. Nat. p. 336.

CYANECULA.

Briss. orn. 3. p. 413.

Partes Europæ boreales incolit pulchra hæc avi-
cula, magnitudine vera in tabula depicta.



R. J. Woodcock Del. et Sculp.

Bill subulate, strait : mandibles nearly equal.
Nostrils obovate.
Tongue jagged or lacerated towards the tip.

SPECIFIC CHARACTER.

Brown MOTACILLA, with the throat blue,
the breast and base of the tail ferruginous,
and the abdomen whitish.

La Gorge-Bleue.

Buff. ois. 5. p. 206.

This elegant species is a native of the northern parts of Europe, and is represented on the plate in its natural size.

PYROSOMA PHOSPHORESCENS.



CHARACTER GENERICUS.

Corpus liberum, oblongo-tubulosum, gelatinosum, altera extremitate apertum.

CHARACTER SPECIFICUS, &c.

PYROSOMA hyalino-virescens, papillosum, ore intus crenato.


PYROSOMA Atlanticum.

Peron. ann. du mus. nation. d'hist. nat.

No. 24. p. 437. t. 72.

De eleganti hoc marino animali primus vere et accurate scripsisse videtur Dominus Peronus, in opere cui titulus “*Annales du Museum National d'Histoire Naturelle.*” Illas incolit maris Atlantici partes quæ circulo æquinoctiali dirimuntur, gregatim radens interdum summas aquas, interdum natans altius submersum. Otiosum et quiescens, ut plurimum sordido-virescit, seu flavescit; simul atque autem sese moverit, fit phosphoro-splendidissimum, singulo contracti corporis actu laminæ si-

millimum ferreæ cum ab igni plene incanduerit ;
singulis autem laxati et extensi vicibus per varios
gradus rubri, aurantii, flavi, leviter cærulei tran-
siens, donec tandem ad colorem naturalem, hyalino
nempe virescentem redierit. Superficiës externa
plurimis papillis seu processibus conicis obsita est :
interna constat e membrana molli et reticulata, sine
ullo visibili intestinorum vestigio. De modo vi-
vendi et sobolem propagandi nihil pro certo com-
pertum est ; probabile tamen est alimentum ei sub-
ministrari eodem modo quo hydræ ; absorpto nempe
succo prædæ prius inclusæ et dein ab ore ejectæ.
Magnitudinem iconis in tabula expressæ sæpius
exsuperat *Pyrosoma phosphorescens*.



THE
PHOSPHORIC PYROSOMA.



GENERIC CHARACTER.

Body nayant, oblong, tubular, gelatinous, open
at one extremity.

SPECIFIC CHARACTER, &c.

Greenish-hyaline, papillose PYROSOMA, with
the mouth crenated within.

PYROSOMA.

Peron annales du Mus. Nat. d'Hist. Nat.
No. 24. p. 437.

This elegant marine animal seems to have been first distinctly described by Mons. Peron, in the work entitled “*Annales du Museum National d'Histoire Naturelle.*” It is a native of the equatorial parts of the Atlantic ocean, where it is sometimes seen in vast numbers, both floating on the surface, and swimming at considerable distances beneath. When in a state of perfect inaction or repose, it is usually

of a dull greenish or yellow cast, and semitransparent; but when in action, is distinguished by a very high degree of phosphoric splendor, exhibiting, during each contraction of its body, the appearance of a bar of iron in the highest degree of incandescence, and, during each relaxation, passing thro' all the gradations of red, orange, yellow, pale-blue, &c. till at length it returns to its first greenish-hyaline aspect. It is externally beset with numerous conical elongations or papillæ, and within appears to be lined merely by a smooth vascular or reticular membrane, without any appearance of intestines, &c. Nothing particular is known relative to its manner of feeding, production of young, &c. It is however probable that it is nourished in the manner of the Polype, viz. by the absorption of the juices of its inclosed prey, which is afterwards ejected by the mouth or opening. In size it often greatly exceeds the figure represented in the annexed plate.

LACERTA LEPIDOPUS.



CHARACTER GENERICUS.

Corpus (plerisque) tetrapodum, elongatum, caudatum, nudum.

CHARACTER SPECIFICUS, &c.

LACERTA olivacea, nigro-punctata, pedibus anterioribus nullis ; posterioribus adactylis, squamosis.

Bipes lepidopodus.

Cepede Ann. Mus. Hist. Nat. No. 21. p. 209.

Quam juxta naturalem magnitudinem expressimus lacertam in Australasia generatur, ab aliis omnibus adhuc cognitis diversa, teste Cepedio, qui illam in opere descripsit quod characteri specifico subjungitur. Pedes habet posticos non divisos in digitos, sed omnino e processibus expansis, orbiculatis, squamosis constantes. Habitu generali accedit ad similitudinem lacertæ quæ *Apus* dicitur, anguinæ, bipedis, &c. ; colore olivaceo seu virescente, punctulis parvulis nigris consperso.



A. J. Nodder Del. J. E. Leach Sculp.

THE
SCALY-FOOTED LIZARD.

GENERIC CHARACTER.

Body (in most species) four-footed, elongated, tailed; without any secondary integument.

SPECIFIC CHARACTER, &c.

Olivaceous LIZARD, speckled with black, without fore-feet, and with scaly, undivided hind-feet.

Le Lepidopode.

Cepede. Ann. Mus. Hist. Nat. No. 21. p. 194.

The Lizard here represented in its natural size is a native of Australasia, and, according to the Count de Cepede, by whom it is described in the work quoted beneath the specific character, differs from every other species yet known in having the hind-feet perfectly destitute of divisions or toes, and consisting merely of a roundish scaly expansion or process. In general habit it resembles the *Lacerta apus*, *anguina*, *bipes*, &c. and is of a greenish or olivaceous colour, variegated by minute blackish specks.

HYALE TRIDENTATA.



CHARACTER GENERICUS.

Animal Clio.

Testa bivalvis, tenuis; valvulis connatis; superiore inflato-convexa, inferiore planiuscula.

CHARACTER SPECIFICUS, &c.

HYALE testa pallida, fusco-flavente, extremitate inferiore tridentata.

ANOMIA tridentata. A.

HYALE.

Cuvier. Ann. Mus. Hist. Nat. No. 21. p. 223.

N. B. Figuræ superiores et inferiores leviter auctæ; cæteræ magnitudine vera exprimuntur.

A testa pellucida nomen adeptum est genus *Hyale*; accurate descriptum et depictum a Domino Cuviero in opere cui titulus "*Annales*," &c. Testam primus detexit Dominus Forskalius, et ad genus *Anomiæ* retulit, cujus habita est species anomala. In mari Mediterraneo enutritum summas


radit aquas quo modo solet genus *Clio*, e Mollus-
corum familia; cui affine admodum est, animal
quod in testa includitur. Longa est testa, ut plu-
rimum, quasi tertiam unciae partem, interdum per-
tingens ultra unciam dimidiatam. Color levissime
fusco-flavescit.

TRIDENTATUS HYAL

TRIDENTATUS HYAL

GENERIC CHARACTER

664



R.M. Sculpt

THE
TRIDENTATED HYALE.



GENERIC CHARACTER.

Animal allied to a Clio.

Shell bivalve, thin, subtransparent; valves connate; the upper convex-inflated, the lower flattish.

SPECIFIC CHARACTER, &c.

Pale yellowish-brown Hyale, with the shell tridentated at the lower part.

HYALE.

Cuvier. Ann. Hist. Nat. No. 21. p. 194.

N. B. The upper and lower figures are slightly magnified; the rest are of the natural size.

The genus *Hyale*, so named from the transparency of the shell, is accurately described and figured by Mons. Cuvier in the "*Annales du M. N. d'Hist. Nat.*" By Forskal, its first describer, it was referred to the genus *Anomia*, of which it was al-

lowed to constitute an anomalous species. It is a native of the Mediterranean sea, and is observed to swim on the surface in the manner of the genus *Clio* among the Mollusca, to which its inhabiting animal is much allied. The usual length of the shell is from one to two-thirds of an inch, and its colour a very pale yellowish brown.

HIRUNDO CAPENSIS.



CHARACTER GENERICUS.

Rostrum minimum, incurvum, subulatum, basi depressum.

Rictus capite amplior.

Lin. Syst. Nat. p. 343.

CHARACTER SPECIFICUS, &c.

HIRUNDO nigro-cærulescens, subtus ochroleuca lineolis fuscis, pileo uropygioque rufis.

HIRUNDO Capensis. H. nigro-cærulescens, subtus flavescens nigricante striata, pileo rufo, rectricibus lateralibus macula alba.

Lath. ind. orn. p. 574.

Juxta veram magnitudinem in tabula depingitur Hirundo Capensis, in Africa australi, ut plurimum, reperta.



R. P. Nodder Del. of F. & S.

1871

THE
CAPE SWALLOW.



GENERIC CHARACTER.

Bill very small, incurved, subulate, depressed
at the base.

Gape wider than the head.

SPECIFIC CHARACTER, &c.

Blueish-black SWALLOW, yellowish white
beneath with small dusky streaks, with the
top of the head and rump rufous.

HIRONDELLE au capuchon roux.

Buff. ois. 6. p. 608.

CAPE SWALLOW.

Lath. Syn. 2. p. 566.

This species of Swallow, which the plate represents in its natural size, is chiefly found in the Southern parts of Africa.

MILLEPORA CERULEA

CHARACTERISTICS

GENERAL DESCRIPTION

Animal Habitat

Geographical Distribution

Depth of Habitat

CHARACTERISTICS

MILLEPORA polymorpha

rules polymorpha

MILLEPORA polymorpha

CHARACTERISTICS

GENERAL DESCRIPTION

Animal Habitat

Geographical Distribution

Depth of Habitat

CHARACTERISTICS

GENERAL DESCRIPTION

Animal Habitat

Geographical Distribution

Depth of Habitat

MILLEPORA CÆRULEA.



CHARACTER GENERICUS.

Animal Hydra.

Corallium poris turbinatis teretibus.

Lin. Syst. Nat. p. 1282.

CHARACTER SPECIFICUS, &c.

MILLEPORA polymorpha subcompressa cærulea, poris inæqualibus.

MILLEPORA plana scabra, laminis crassis varie tortuosis subdivisa, apicibus sæpe lobatis porisque substellatis cylindricis utrinque instructis.

Soland. et Ell. zooph. p. 142. t. 12. et 56.

CORALLOIDES philippensis cærulea.

Pet. gaz. t. 10. f. 12.

In maribus Indicis præcipue conspicitur formosum hoc corallium, magnitudine vera in tabula expressum.



BLUE MILLEPORE.



GENERIC CHARACTER.

Animal allied to a Polype.

Coral with very numerous cylindric pores.

SPECIFIC CHARACTER, &c.

Polymorphous, subcompressed, blue Millepore,
with unequal pores.

Blue MILLEPORE.

Soland. et Ellis zooph. t. 12. et 56.

Blue INDIAN CORAL.

This beautiful Coral is chiefly seen in the Indian seas, and is represented in its natural size on the annexed plate.



SPHINX LABRUSCÆ.



CHARACTER GENERICUS.

Antennæ medio crassiores, seu utraque extremitate attenuatæ, subprismaticæ.

Alæ deflexæ, (volatu graviore vespertino seu matutino.)

Lin. Syst. Nat. p. 796.

CHARACTER SPECIFICUS, &c.

SPHINX olivacea, alis integris, posticis subcæruleis nigro-fasciatis, margine interiore sanguineis.

SPHINX LABRUSCÆ. S. alis subfuscis; primoribus subtus puncto albo, abdomine lateribus punctis quinque albis.

Lin. Syst. Nat. p. 800.

Insectum depinximus elegantissimum, in America australi generatum, a Domina Meriana primo, ut videtur, descriptum et expressum in splendido opere quod inscribitur "Metamorphoses insectorum

Surinamensium." Larva, cui præ alio cibo gratae
sunt variae vitium species, teste Meriana, Augusto
mense in chrysalidem convertitur, e qua Septembri
insequente erumpere solet ipsa Sphinx.



H. P. Nodder. Del. & Sculp.

THE
WILD-VINE SPHINX.



GENERIC CHARACTER.

Antennæ subprismatic, thickest in the middle;
and attenuated at each extremity.

Wings deflected.

SPECIFIC CHARACTER, &c.

Olivaceous SPHINX, with entire wings; the
lower ones blueish with black bars and
blood-red interior margin.

MERIAN. *Ins. Surin.* t. 34.

The highly elegant insect here represented is a native of South-America, and seems to have been first described and figured by the celebrated Madam Merian, in her splendid work on the insects of Surinam. Its caterpillar feeds principally on the different species of Vine. According to Madam Merian it changes to a chrysalis in the month of August, from which in that of September emerges the Sphinx.

feriores autem albido-flavæ. Cauda subforcipata, tres pollices longa; rectrices duodecim, fuscæ, marginibus albido-flavicantibus.

“Femina differt coloribus capitis et trunci; illa nimirum supra tota a fronte ad caudam usque sordide olivaceo-ferruginea fusco maculata, subtus tota ex albido flava.

“Habitat in submontanis promontorii utriusque, et septentrionalis et meridionalis Caucasi, circa thermas ad fluvium Terek obvias et in Georgia circa Tessium.”

669



R. P. Nodder. Del. et Sculp.

BLACK-CROWNED TANAGER.

GENERIC CHARACTER.

Bill conic, acuminated, a little inclining towards the point; the upper mandible slightly ridged, and notched near the end.

SPECIFIC CHARACTER, &c.

Ferruginous-brown TANAGER, yellow beneath, with black crown and cheeks, and wings streaked with white.

Black-crowned TANAGER.

Lath. Syn. 2. p. 223.

This species is described by Guldenstadt in the Petersburg Transactions. Its size is that of the common Bunting: the crown and cheeks are black, the neck and back brown-ferruginous, the rump yellowish-ferruginous, and the whole bird beneath deep yellow; the wings which, when closed, reach to the middle of the tail, are streaked longitudinally with white; the tail is slightly forked, measuring about three inches in length. The female differs in being entirely of a dull olive-ferruginous above, spotted with black, and beneath yellowish white.

It is an inhabitant of the Caucasian mountains and of the country of Georgia.

SPHINX PANOPUS.

CHARACTER GENERICUS.

Antennæ medio crassiores, seu utraque extremitate attenuatæ, subprismaticæ.

Alæ deflexæ (volatu graviore respertino seu matutino).

Lin. Syst. Nat. p. 796.

CHARACTER SPECIFICUS, &c.

SPHINX alis gilvis fusco fasciatis, primoribus ocello nigro ad angulum interiorem.

SPHINX PANOPUS.

Cram. t. 224. f. A. B.

Insulam Javam incolet Sphinx Panopus, magnitudine vera in tabula depictus.





RPN Del. et Sculp.

PANOPUS.

GENERIC CHARACTER.

Antennæ subprismatic, thickest in the middle,
and attenuated at each extremity.

Wings deflected.

SPECIFIC CHARACTER, &c.

Sphinx with buff-coloured wings barred with
brown ; the upper pair marked by a black
ocellated spot at the interior angle.

PANOPUS.

Cram. pap. exot. t. 224. f. A. B.

This insect is a native of the island of Java, and is
represented on the plate in its natural size.

GORGONIA CERATOPHYTA.

CHARACTER GENERICUS.

Animal crescens plantæ facie.

Hydræ sparsæ e poris lateralibus.

Stirps radicata, cornea, continua, ramosa, basi explanata, cortice obducta.

CHARACTER SPECIFICUS, &c.

GORGONIA subdichotoma purpurea, ramis virgatis, poris bifariis.

GORGONIA subdichotoma, axillis divaricatis, ramis virgatis bisulcatis, cortice rubro, poris bifariis.

Lin. Syst. Nat. Gmel. p. 3800.


GORGONIA CERATOPHYTA.

Soland. et Ellis Zooph. p. 81. t. 12.

Corallina fruticosa purpurea.

C. Bauh. pin. 366.

Circa oras Americanas præcipue conspicitur formosum hoc Corallium, altitudine, ut plurimum, pedali vel sesquipedali.



PURPLE GORGONIA.

GENERIC CHARACTER.

Animal growing with the habit of a plant.

Polypes scattered from the lateral pores.

Stem fixed, horny branched, covered with a soft bark.

SPECIFIC CHARACTER, &c.

Subdichotomous purple GORGONIA, with twiggy branches, and double rows of white polypes.

Purple shrubby GORGONIA.

This beautiful coral is principally seen about the American coasts : its general height is about a foot or eighteen inches.

LEPAS AURITA.

CHARACTER GENERICUS.

Animal Triton.

Testa multivalvis, inæquivalvis, basi affixa.

Lin. Syst. Nat. p. 1107.

CHARACTER SPECIFICUS, &c.

LEPAS pedunculata fusca membranacea, appendicibus auriculatis.

LEPAS testa membranacea ventricosa tubo insidente, ore octovalvi dentato, tubulo gemino aurita.

Lin. Syst. Nat. p. 1110.

Speciem quam depinximus, marium Arcticorum incolam, primum descripsisse arbitramur Ellisium in Actis Anglicis. Adhæret, ut solet ferme reliquum genus, plurimis marinis substantiis. Quas ostendit tabula Lepadas repræsentavimus grandiori cuidam speciei affixas, cui nomen inditum est corporis figuræ conveniens ; Lepas nempe *Diadema*.



THE
EARED BARNACLE.

GENERIC CHARACTER.

Animal a Triton.

Shell multivalve, with unequal valves, affixed
by the base.

SPECIFIC CHARACTER, &c.

Brown membranaceous pedunculated BARNACLE, with auriculated appendages.

Auriculated BARNACLE.

This species is a native of the northern ocean, and seems to have been described by Mr. Ellis in the Philosophical Transactions. Like others of this genus, it adheres to various marine substances; and the individuals on the present plate are represented adhering to a larger species, called from its figure *Lepas Diadema*.

PSITTACUS UNDULATUS.

CHARACTER GENERICUS.

Rostrum aduncum, mandibula superiore mobili,
cera instructa.

Nares in rostri basi.

Lingua carnosae, obtusa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

CHARACTER SPECIFICUS, &c.

PSITTACUS macrourus viridis, supra fusco
undulatus, gula flavente cœruleo maculata,
rectricibus basi flavis.

Ostendit tabula perpulchram speciem psittacinam, Australasiæ incolam, quamque censeo non antea fuisse descriptam. Superiora avis a rostro nempe ad uropygium flavo-viridi-pallent, undulis plurimis fuscis et linearibus transverse decorata, gradatim majoribus prope humeros et imum dorsum. Fuscae sunt alarum pennae, marginibus olivaceo-subflavescentibus. Inferiora cum uropygio virent eleganter pallida. Gula flavo-pallet, maculis paucis cæruleis, lunulisque nigris, huc illuc utrinque conspersa. Cauda cuneata cyanea est, fascia flavissima per omnes pennas in obliquum ducta, exceptis duabus mediis, quæ cæteras longitudine exsuperant. Rostrum pedesque fusca. Exprimitur in tabula vera avis magnitudo.



Dr. S. Del.

R. Madder. Sculp.

THE
UNDULATED PARRAKEET.

GENERIC CHARACTER.

Bill hooked : upper mandible moveable.
Nostrils round, placed in the base of the bill.
Tongue fleshy, broad, blunt at the end.
Legs short : feet scansorial.

SPECIFIC CHARACTER, &c.

Long-tailed green PARRAKEET, undulated
above with brown ; the throat yellowish,
with blue spots, and the tail-feathers yellow
at the base.

The highly elegant species of Parrakeet represented on the present plate in its natural size, is an inhabitant of New Holland, and seems to have been hitherto undescribed. The upper parts of the bird, from the bill to the rump, are of a pale yellowish green, beautifully crossed by numerous linear brown undulations, which become gradually larger as they approach the back and shoulders ; the wing-feathers are brown, with pale olive-yellow edges ; the under parts of the bird, together with the rump, are of an elegant pale green ; the throat pale yellow, mottled on each side with a few small deep blue scattered spots, accompanied by

small black crescents: the tail is of a cuneated form, and of a deep-blue colour, with a bright yellow bar running obliquely across all the feathers except the two middle ones, which considerably exceed the rest in length: the bill and legs are brown.

GORGONIA CRASSA ?

CHARACTER GENERICUS.

Animal crescens plantæ facie.

Hydræ sparsæ e poris lateralibus.

Stirps radicata, cornea, continua, ramosa, basi explanata, cortice obducta.

CHARACTER SPECIFICUS, &c.

GORGONIA teres dichotoma, ramis crassis adscendentibus, carne violacea.

GORGONIA CRASSA ? G. dichotoma, &c. &c.

Soland. & Ellis Zooph. p. 91.

GORGONIA CRASSA ?

Lin. Syst. Nat. Gmel. p. 3806.

Super rupes maris Indici et Americani conspicitur Gorgonia crassa, cujus veram magnitudinem exprimit tabula.



THICK GORGONIA.

GENERIC CHARACTER.

Animal growing with the habit of a plant.

Polypes scattered from the lateral pores.

Stem fixed, horny, branched, covered with a soft bark.

SPECIFIC CHARACTER, &c.

Round-stemmed, dichotomous GORGONIA,
with thick, ascendant branches, and violet-coloured bark.

This species of Gorgonia is principally seen on the rocks of the Indian and American seas, and is represented in its natural size.

OSTREA DENTICULATA.

CHARACTER GENERICUS.

Animal Tethys.

Testa bivalvis, inæquivalvis, subaurita.

Cardo edentulus, fossula cava ovata, striisque lateralibus transversis.

Lin. Syst. Nat. p. 1144.

CHARACTER SPECIFICUS, &c.

OSTREA testa rotunda, foliacea, valva altera plana parasitica, cardine utrinque denticulato.

Born. test. Mus. Cæs. Vind. p. 113.
t. 6. f. 9, 10.

OSTREA major sulcata, inæqualiter utrinque ad cardinem denticulata.

List. Conch. t. 193, 194.

Circa promontorium bonæ spei, scopulis affixa, conspicitur Ostrea denticulata, interdum major quam in tabula ostenditur.



THE
DENTICULATED OYSTER.

GENERIC CHARACTER.

Animal resembling a Tethys.

Shell bivalve, subauriculated.

Hinge generally toothless, with an ovate or sub-trigonal fossule, and lateral transverse streaks.

SPECIFIC CHARACTER, &c.

OYSTER with foliaceous, rounded shell; one valve flat and parasitic, with the hinge denticulated on each side.

Born. Test. Mus. Cæs. Vindob. t. 6. f. 9, 10.

The denticulated Oyster is found adhering to rocks about the Cape of Good Hope, and is sometimes larger than the specimen represented on the annexed plate.

MYRMELEON GRANDE.

CHARACTER GENERICUS.

Os maxillosum, dentibus duobus. Palpi quatuor elongati.

Stemmata nulla.

Cauda maris forcipe e filamentis duobus rectiusculis.

Antennæ clavatæ, longitudine thoracis.

Alæ deflexæ.

Lin. Syst. Nat. p. 913.

CHARACTER SPECIFICUS, &c.

MYRMELEON antennis gracilibus clavatis, alis fusco maculato-fasciatis.

Drur. Ins. Exot. 3. t. 41.

Disseruere interdum physici de eleganti hoc insecto, quasi mera esset varietas Myrmeleontis *libelluloidis*. Rectius fortasse statuatur speciem esse revera diversam. In Africa generatur Myrmeleon grande, et veram ejus magnitudinem cernere est in tabula.



THE
GREAT MYRMELEON.

GENERIC CHARACTER.

Mouth furnished with jaws and two teeth.

Feelers four, elongated.

Stemmata none.

Tail (of the male) forcipated by two straitish filaments.

Antennæ equalling the thorax in length, and clavated.

Wings deflected.

SPECIFIC CHARACTER, &c.

MYRMELEON with slender, clavated antennæ, and wings barred by patches of brown.

Drury Ins. 3. t. 41.

This elegant insect, which is sometimes regarded as a variety only of the Myrmeleon libelluloides, may, perhaps, with greater propriety be considered as constituting a distinct species. It is a native of Africa, and is represented in its natural size.

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

THE GREAT BRITISH

LOXIA LUDOVICIANA.

CHARACTER GENERICUS.

Rostrum conico-gibbum, frontis basi rotundatum versus caput; mandibula inferiore margine laterali inflexa.

Nares in basi rostri.

Lingua integra.

Lin. Syst. Nat. p. 299.

CHARACTER SPECIFICUS, &c.

LOXIA nigra, subtus alba, pectore roseo, alis fascia duplici alba.

LOXIA LUDOVICIANA. L, nigra, pectore, ventre, alarum fascia, basique remigum albis.

Lin. Syst. Nat. p. 306.

Coccothraustes LUDOVICIANA.

Briss. av. 3. p. 247. t. 12. f. 2.

Americæ septentrionalis varias regiones incolit Loxia Ludoviciana, cujus veram magnitudinem ostendit tabula.



R.R. Noddy. Del. et. Sculp.

THE
LUDOVICIAN GROSSBEAK.

GENERIC CHAPACTER.

Bill, strong, thick, and convex.

Nostrils small.

Tongue truncated.

SPECIFIC CHARACTER, &c.

Black GROSSBEAK, white beneath, with
rose-coloured breast, and a double white
bar on the wings.

Red-breasted GROSSBEAK.

Lath. Syn. 3. p. 126.

Penn. Arch. Zool. p. 350.

Le Rose-gorge.

Buff. Ois. 3. p. 460.

GROS-BEC de la LOUISIANE.

Pl. Eul. 153. f. 2.

The Ludovician Grossbeak is a native of several
parts of North America, and is exhibited on the plate
in its natural size.

NERITA POLITA
AND
NERITA PENNATA.

CHARACTER GENERICUS.

Animal Limax.

Testa univalvis, spiralis, gibba, subtus planiuscula.

Apertura semiorbicularis, labio columellæ transverso, truncato, planiusculo.

Lin. Syst. Nat. p. 1251.

CHARACTER SPECIFICUS, &c.

NERITA POLITA. N. testa lævi, vertice obliterato, labiis utrisque dentatis.

Lin. Syst. Nat. p. 1254.

NERITA POLITA. N. testa lævi obtusa, labio dentato, labro crenulato.

Born. Mus. Cæs. p. 405. t. 17. f. 15, 16.

NERITA PENNATA. N. testa olivacea albo variegata, maculis transversis pennatis nigris.

NERITA PENNATA. N. testa rotundata lævi, labio plano crenulato.

Born. Mus. Cæs. p. 404. t. 17. f. 11, 12.

Si generis Neritæ formam generalem spectes, quæ bella admodum et concinna est, videbitur propius ac-

cedere ad similitudinem generis Helicis. Sed quoniam
major pars Neritarum maria incolat, datæ sunt iis
testæ densiores multo et validiores quam helicibus vel
terrestribus vel fluviatilibus. In mari Indico generan-
tur ambæ species in tabula depictæ.



W. B. D. & Co. London

The figure of the nautilus is produced by a single nautilus of a species. In point of form it bears a great resemblance to that of the nautilus, but as by the the greatest number of species are natives of the sea they are of a much thicker and stronger nature than either the land or fresh-water nautilus. Both the species here represented are produced in the Indian sea.

THE
GLOSSY NERITE
AND
FEATHERED NERITE.

GENERIC CHARACTER.

Animal allied to a Limax or Slug.

Shell univalve, spiral, gibbose, flattish beneath.

Aperture semiorbicular, with the lip of the pillar transverse, truncated, and flattish.

SPECIFIC CHARACTER, &c.


GLOSSY NERITE. Smooth, whitish Nerite, with reddish and dusky variegations and streaks.

Born. Mus. Cæs. pl. 17. f. 15, 16.

FEATHERED NERITE. Olivaceous Nerite, with white variegations, and transverse, feather-shaped, black spots.

Born. Mus. Cæs. pl. 17. f. 11, 12.

The genus *Nerita* is possessed of a singular neatness of appearance. In point of form, it bears a near resemblance to that of *Helix*, or Snail; but as by far the greater number of species are natives of the sea, they are of a much thicker and stronger fabric than either the land or fresh-water snails. Both the species here represented are produced in the Indian seas.



SPHINX JATROPHÆ.

CHARACTER GENERICUS.

Alæ subprismaticæ, utroque fine attenuatæ.

Lingua exserta (plerisque).

Palpi duo reflexi.

Alæ deflexæ.

Lin. Syst. Nat. Gmel. p. 2371.

CHARACTER SPECIFICUS, &c.

SPHINX grisea, alis superioribus nigro undulatis, corpore utrinque ocellis tribus luteis nigro marginatis.

SPHINX JATROPHÆ. S. alis subdentatis, posticis nigris basi rufis fasciaque fenestrata, capite bicorni.

Fab. Sp. Ins. 2. p. 143.

In Surinamia innascitur Sphinx Jatrophæ, nec non in aliis Americæ Australis regionibus. Memoravit et depinxit hanc speciem celeberrima Domina Merian. Larva Jatrophæ gossypifoliæ ramulos præcipue depascitur.



THE
JATROPHA SPHINX.

GENERIC CHARACTER.

Antennæ subprismatic, attenuated at each extremity.

Tongue generally exerted.

Feelers two, reflex.

Wings deflected.

SPECIFIC CHARACTER, &c.

Grey SPHINX, with the upper wings undulated with black, and the body ocellated on each side by three yellow spots with black margins.

Merian Ins. Surin. pl. 38.

This insect is an inhabitant of Surinam, as well as of some other parts of South America, and is figured by the celebrated Madam Merian, in her work on the Insects of Surinam. The caterpillar feeds chiefly on the shoots of the *Jatropha gossypifolia*.

ALCEDO TRIBRACHYS.

CHARACTER GENERICUS.

Rostrum trigonum, crassum, rectum, longum.

Lingua carnosae, brevissima, plana, acuta.


Pedes gressorii plerisque.

Lin. Syst. Nat. p. 178.

CHARACTER SPECIFICUS, &c.

ALCEDO cyanea, subtus ferruginea, alis nigricantibus, pedibus tridactylis.

Australasiam incolit Alcedo Tibrachys. Ostendit tabula veram avis magnitudinem, a nemine, ut opinor, antea descriptæ aut depictæ.





THE
TRIDIGITATED KINGFISHER.

GENERIC CHARACTER.

Bill trigonal, thick, straight, long.

Tongue fleshy, very short, flat, sharp-pointed.

Feet gressorial.

SPECIFIC CHARACTER, &c.

Deep-blue KINGSFISHER, ferruginous beneath, with blackish wings and tridactyle feet.

The present species, which is a native of Australasia, does not appear to have been hitherto either figured or described. It is represented on the plate in its natural size.

PERCA? MACULATA.

CHARACTER GENERICUS.

Squamæ duræ, asperæ.

Opercula spinosa.

Bloch. ichth. 2. p. 56.

CHARACTER SPECIFICUS, &c.

PERCA? albo-flavescens, rubro guttata.

PERCA MACULATA. P. alba, guttis rubris.

Bloch. t. 313.

Sparus Atlanticus?

Gen. Zool. 4. p. 453.

In mari Atlantico præcipue repertus, sedem sibi jure vindicat hic piscis inter pulcherrimos sui generis. Colore interdum variat, et in longitudinem quasi quindecim vel octodecim unciarum crescit.



R.P. Woodw. Del. et Sculp.

RED-SPOTTED PERCH?

GENERIC CHARACTER.

Gill-covers spiny.

Scales hard and rough.


Bloch.

SPECIFIC CHARACTER, &c.

Yellowish-white PERCH, with red spots.

Red-spotted American PERCH.

This fish is principally seen in the American seas, and may justly be considered as one of the most beautiful of its genus. In colour it sometimes varies, and grows to the length of fifteen or eighteen inches.



HELIX CITRINA.

CHARACTER GENERICUS.

Animal Limax.

Testa univalvis, spiralis, subdiaphana, fragilis.

Apertura coarctata, intus lunata seu subrotunda,
segmento circuli dempto.

CHARACTER SPECIFICUS, &c.

HELIX testa subumbilicata convexa obtusa flavescens, fascia fusca.



Lin. Syst. Nat. p. 1245.

Gualt. test. t. 3. f. D. E.

Seb. mus. 3. t. 39. f. 1, 2.

Born. mus. Cæs. Vind. test. t. 13. f. 14, 15.

Varias Asiæ atque Americæ regiones incolit Cochlea hæc elegantissima, magnitudine vera in tabula depicta.



90 No 10 Dec Del 70 24/1/71

THE
CITRON SNAIL.


GENERIC CHARACTER.

Animal resembling a Slug.
Shell spiral sub-pellucid,
Aperture semilunar.

SPECIFIC CHARACTER, &c.

SNAIL with citron-coloured, subumbilicated,
convex, obtuse shell, with a dusky band.
The citron-coloured Jamaica SNAIL.
The yellow Indian SNAIL.

This highly elegant shell is a native of many parts of
Asia and America, and is represented on the plate
in its natural size.



LITTON SNAIL

CHARACTERISTICS

CHARACTERISTICS

Shells resembling a flag.

Shells spiral and polished.

Shells containing...

CHARACTERISTICS

SNAIL, with a smooth, polished, and polished.

conver, oblate, with a dark band.

The citron-colored Japanese SNAIL.

The yellow Italian SNAIL.

The light elegant shell is a native of the...

and the African, and is represented on the plate.

In its natural state...

GORGONIA BRIAREUS.

CHARACTER GENERICUS.

Animal crescens plantæ facie.

Hydræ sparsæ e poris lateralibus.

Stirps radicata, cornea, continua, ramosa, basi explanata, cortice obducta.

CHARACTER SPECIFICUS, &c.

GORGONIA teres, crassa, cinerea, basi explanata, osse vitreo-aciculato, hydri magnis cirratis.

GORGONIA BRIAREUS. G. subramosa teres crassa, basi supra rupes late explanata, carne interna subalbida, externe cinerea, polypis majoribus octotentaculatis cirratis, osse ex aciculis vitreis purpureis, inordinate sed longitudinaliter compactis composito.

Soland. et Ellis Zooph. p. 93. t. 14. f. 1.

Circa oras Americanas præcipue conspicitur Gorgonia Briareus, altitudine decem vel quindecim uncias æquans. Colore interdum variat.

THE
THICK-ARMED GORGONIA.

GENERIC CHARACTER.

Animal growing with the habit of a plant.

Polypes scattered from the lateral pores.

Stem fixed, horny, branched, covered with a soft bark.

SPECIFIC CHARACTER, &c.

Round-armed, thick, cinereous GORGONIA,
with expanded base, glassy-fibred bone,
and large cirrated polypes.

The GORGON BRIAREUS.

Soland. et Ellis Zooph. p. 93. pl. 14. f. 1.

This species of Gorgonia is chiefly found about the American coasts, and grows to the height of ten or fifteen inches. In colour it sometimes varies.



INDEX.

PL.

- 638. *Aplysia punctata*.
- 681. *Alcedo tibrachys*.
- 642. *Cancer Mantis*.
- 655. ——— *Scyllarus*.
- 648. { *Cypræa fragilis*.
- *Arabica*.
- 658. *Corallina Flabellum*.
- 671. *Gorgonia ceratophyta*.
- 674. ——— *crassa*.
- 684. ——— *Briareus*.
- 683. *Helix citrina*.
- 664. *Hyale tridentata*.
- 665. *Hirundo Capensis*.
- 637. *Lanius gutturalis*.
- 645. ——— *jocosus*.
- 672. *Lepas aurita*.
- 663. *Lacerta lepidopus*.
- 677. *Loxia Ludoviciana*.
- 647. *Lucernaria quadriloba*.
- 643. *Medusa Nummularia*.
- 649. *Motacilla Rossica*.
- 661. ——— *Suecica*.
- 666. *Millepora cærulea*.
- 676. *Myrmeleon grande*.
- 678. *Nerita polita et pennata*.
- 675. *Ostrea denticulata*.
- 639. ——— *Jacobæa*.
- 641. *Pavo spicifer*.
- 640. *Phalæna Apollonia*.
- 646. ——— *Tarquinia*.
- 644. ——— *Fabia*.
- 656. ——— *Janus*.
- 659. ——— *Paphia, larva*.
- 660. ——— *Paphia*.
- 650. *Papilio Amphrysus*.
- 653. *Psittacus melanotus*.
- 657. ——— *zonarius*.
- 673. ——— *undulatus*.
- 682. *Perca maculata*.
- 662. *Pyrosoma phosphorescens*.
- 651. *Spondylus gædaropus*.
- 652. *Sparus melanopterus*.
- 667. *Sphinx labruscæ. larva*.
- 668. ——— *labruscæ*.
- 670. ——— *Panopus*.
- 679. ——— *Jatropha. larva*.
- 680. ——— *Jatropha*.
- 669. *Tanagra melaniçtera*.

PL.

- 638. *Aplysia speckled*.
- 637. *Butcher-Bird red-throated*.
- 650. *Butterfly Amphrysus*.
- 672. *Barnacle eared*.
- 642. *Crab long-bodied*.
- 655. *Crab oval-tailed*.
- 660. *Coralline fan*.
- 648. *Cowry brittle and Arabian*.
- 671. *Gorgonia purple*.
- 674. ——— *thick*.
- 684. ——— *thick-armed*.
- 677. *Grossbeak Ludovician*.
- 664. *Hyale tridentated*.
- 681. *Kingfisher tridigitated*.
- 647. *Lucernaria four-lobed*.
- 663. *Lizard scaly-footed*.
- 643. *Medusa nummular*.
- 644. *Moth Fabia*.
- 640. ——— *Apollonia*.
- 646. ——— *Tarquinia*.
- 656. ——— *Janus*.
- 658. ——— *Paphia-larva*.
- 659. ——— *Paphia*.
- 654. *Mackrel Plumerian*.
- 666. *Millepore blue*.
- 676. *Myrmeleon great*.
- 678. *Nerite glossy and feathered*.
- 675. *Oyster denticulated*.
- 653. *Parrakeet black-backed*.
- 673. ——— *undulated*.
- 657. *Parrot zoned*.
- 662. *Pyrosoma phosphoric*.
- 641. *Peacock spike-crested*.
- 682. *Perch red-spotted*.
- 661. *Red-Breast Swedish*.
- 639. *Scallop Jacobæan*.
- 683. *Snail citron*.
- 645. *Shrike red-vented*.
- 665. *Swallow Cape*.
- 652. *Sparus black-finned*.
- 651. *Spondylus subauriculated*.
- 667. *Sphinx wild vine. larva*.
- 668. ——— *wild vine*.
- 679. ——— *Jatropha. larva*.
- 680. ——— *Jatropha*.
- 670. ——— *Panopus*.
- 669. *Tanager black-crowned*.
- 649. *Warbler Russian*.



